

Collective Excitations in Quantum Hall Liquid Crystals: Single-Mode Approximation Calculations

Cintia M. Lapilli and Carlos Wexler

Department of Physics and Astronomy, University of Missouri–Columbia, Columbia, Missouri 65211, USA
(Dated: August 19, 2005, new version: December 30, 2005)

A variety of recent experiments probing the low-temperature transport properties of quantum Hall systems have suggested an interpretation in terms of liquid crystalline mesophases dubbed *quantum Hall liquid crystals*. The single mode approximation (SMA) has been a useful tool for the determination of the excitation spectra of various systems such as phonons in ^4He and in the fractional quantum Hall effect. In this paper we calculate (via the SMA) the spectrum of collective excitations in a quantum Hall liquid crystal by considering *nematic*, *tetratic*, and *hexatic* generalizations of Laughlin's trial wave function having two-, four- and six-fold broken rotational symmetry, respectively. In the limit of zero wavevector $\mathbf{q}$ the dispersion of these modes is singular, with a gap that is dependent on the direction along which $\mathbf{q} = 0$ is approached for *nematic* and *tetratic* liquid crystalline states, but remains regular in the *hexatic* state, as permitted by the fourth order wavevector dependence of the (projected) oscillator strength and static structure factor.

PACS numbers: 73.43.-f, 73.20.Mf, 64.70.Md, 52.27.Aj

I. INTRODUCTION

For more than two decades two-dimensional electron systems (2DES), and in particular the quantum Hall effect (QHE)^{1,2,3} have been constant sources of complex and unexpected behavior, perhaps with no equal in the realms of condensed matter physics. The unique combination of extremely high mobilities ($\mu \sim 10^7$ m/Vs) in GaAs/Al_xGa_{1-x}As heterostructures, low temperatures ($T < 100$ mK), enhancement of interactions due to the reduced dimensionality, and relative quenching of the kinetic energy in strong magnetic fields due to Landau level (LL) quantization, has allowed the emergence of complex and striking behavior due to subtle correlation effects. In fact, from the discovery of the integer and fractional QHE's in the early eighties¹ (leading to the award of two Nobel prizes), to the existence of novel fractionally charged² and composite particles⁴ and many newer interesting many-body phenomena, the QHE has been a consistently active and exciting area of research.

The physics of 2DES in partially filled LL's is inherently complex due to the high degeneracy of the "unperturbed" ground state (i.e. without the Coulomb interaction). The first successful theoretical approach to this system was proposed shortly after the discovery of the FQHE¹ by Laughlin, who proposed his famous trial-wavefunction²

$$\Psi_{1/m} = \prod_{i < j}^N (z_i - z_j)^m e^{-\frac{1}{4} \sum_{k=1}^N |z_k|^2}, \quad (1)$$

to describe states at filling factor $\nu = 1/m$, with m an odd integer. Here $z_j = x_j + iy_j$ is the position of the j^{th} electron in the complex plane and we work in units of the magnetic length $l_0 = [\hbar/eB]^{1/2}$. The "goodness" of Laughlin state originates in that the nodal hyper-surfaces on which the many-body wavefunction vanishes coincide with the ones in which the particles are in contact. Whereas the vanishing of the wavefunction when two (spin aligned) electrons are in contact is required by Fermi statistics, Laughlin state has multiple nodes at those points, thus reducing the Coulomb repulsion.

Later, Jain proposed⁴ the elegant conceptual framework of the composite fermions (CF) which unifies all hierarchies of integer and fractional QHE's, along with the intermediate states in between QH plateaus: an even number $2p$ of vortices is attached to each electron, also lowering the Coulomb energy, and (in a "mean field" approximation) the "magnetic fluxes" associated with the vortices lead to a reduction of the effective magnetic field: $B^* = B - 2p\Phi_0 n_e$ ($\Phi_0 = h/e$ is the flux quantum and n_e is the electron density), resulting in an effective filling factor $\nu^* = \nu/(1 - 2p\nu)$. It is easy to see that for the strongest FQHE states ν^* is an integer (e.g. for $\nu = 1/3$ and $p = 1$ $\nu^* = 1$), leading to the interpretation of the FQHE of electrons as a simple IQHE of the CF's; whereas the intermediate regions (e.g. $\nu = 1/2, 1/4$) correspond to $\nu^* \rightarrow \infty$, i.e. a vanishing effective magnetic field: $B^* = 0$.

Since 1999, magnetotransport experiments have uncovered a variety of surprising results at low temperatures ($T \lesssim 100$ mK), for example: extreme anisotropies⁵ and apparent competition between different ordered phases⁶ in the intermediate regions between quantum Hall plateaus at high LL's, the melting transition between a Wigner crystal (WC) at $\nu \simeq 1/7$,⁷ and the microwave conductivity evidence of structural phase transitions in partially filled LL's.⁸ A large body of evidence corresponding to seemingly distinct phenomena may be partly understood in terms of a simple conceptual point: that the many-body states involved have an intrinsic crystalline or liquid crystalline order,^{9,10,11,12,13,14,15,16,17} be it *smectic*,^{9,11} *nematic*,^{10,11,12,13} *tetratic* or *hexatic*.¹³

Generalizations of Laughlin wavefunction [Eq. (1)] with discrete broken rotational symmetry (BRS) have been proposed in the past^{12,13,18} as candidates for nematic or hexatic states^{12,13} in order to understand anisotropic transport observed in the intermediate regions,⁵ or the melting of the WC at $\nu = 1/7$.⁷ In fact, the motivation for these states arises from the fact that it is generally expected that melting in 2D may occur through a topological Kosterlitz-Thouless-type (KT) transition.¹⁹ For 2D melting, the reliable Kosterlitz-Thouless-Halperin-Nelson-Young (KTHNY) theory^{19,20} predicts that, in fact, an *intermediate liquid crystalline phase* will exist be-

tween a solid and a liquid phase, which will exhibit no *translational* order, and only a quasi-long-range order for the orientational order below the KT transition temperature. These arguments were used by Wexler and Dorsey¹¹ to calculate qualitatively correct anisotropic-isotropic transition temperatures for the quantum Hall liquid crystal in the transitional regions at high LL's.⁵

In this paper we consider the spectrum of collective excitations for a family of liquid crystalline states in a partially filled LL. These states are generated so as to satisfy the following criteria which we consider reasonable for understanding the dynamics: (i) states must obey Fermi statistics, i.e., they must be *odd* under the exchange of any pair of electrons; (ii) the states must be translationally invariant (far enough from boundaries); (iii) there must be a broken rotational symmetry belonging to the proper symmetry group (i.e., C_2 for a *nematic*, C_4 for a *tetragic*, and C_6 for a *hexatic*; additional symmetries are possible in principle, e.g. with a C_{10} symmetry, we have not explored such possibilities); (iv) states and excitations must reside entirely in the LLL to avoid the large cyclotron energy cost $\hbar\omega_c$. Note, as we will show later, that the properties of *any* excited LL may be readily obtained from the properties of the LLL.

States that satisfy the abovementioned requirements have been proposed and studied in detail^{12,13,18} for filling factors $\nu = 1/3$, $1/5$, and $1/7$. These are found by splitting the “extra” vortices of the Laughlin (or other CF) states around the electron, while obeying the required symmetries:

$$\Psi_{1/(2p+1)}^\alpha = \left\{ \prod_{i < j}^N \left[\prod_{\mu=1}^{2p} (z_i - z_j - \alpha_\mu) \right] \right\} \times \prod_{i < j}^N (z_i - z_j) e^{-\frac{1}{4} \sum_{k=1}^N |z_k|^2}, \quad (2)$$

where the complex numbers α_μ are distributed in pairs of opposite value in the complex plane (to satisfy Fermi statistics). For the states with the highest discrete symmetry at each filling factor we may take

$$\alpha_\mu = \alpha e^{i2\pi(\mu-1)/2p}, \quad \mu \in \{1, 2, \dots, 2p\}, \quad (3)$$

and without loss of generality α can be taken to be real. The wavefunction in Eq. (2) represents a homogeneous liquid crystalline state at $\nu = 1/(2p+1)$, is anti-symmetric, lies entirely in the LLL, and is smoothly connected to the isotropic Laughlin state for $\alpha = 0$. Figure 1 depicts the nodal distribution for various states of Eq. (2). A remarkable feature of these states is that they possess an underlying charge density wave (CDW), but these CDWs are melted by fluctuations, and overall the system is translationally invariant.¹³

II. THE SINGLE MODE APPROXIMATION

To calculate the excitation spectrum we use the single mode approximation (SMA),^{21,22} which reliably provides the first moment (mean) of the energy of the excitations (for a given




FIG. 1: Nodal distribution for $z_{ij} \equiv z_i - z_j$ for a quantum Hall *nematic* at $\nu = 1/3$, *tetragic* at $\nu = 1/5$, and *hexatic* at $\nu = 1/7$.

wavevector $\mathbf{k}$) that are coupled to the ground state by means of the density operator.^{21,22,23} The SMA was first used by Feynman in 1953 to accurately calculate the spectrum of phonons in superfluid ^4He .²¹ The essence of the method originates on the assumption that the ground state of a system of bosons has a scarcity of long-wavelength excitations. Under those circumstances, the variational wavefunction for an excitation corresponding to a density-wave can be written as

$$\phi_{\mathbf{k}}(\mathbf{r}_1, \dots, \mathbf{r}_N) = N^{-1/2} \rho_{\mathbf{k}} \psi_0(\mathbf{r}_1, \dots, \mathbf{r}_N), \quad (4)$$

where $\rho_{\mathbf{k}} = \sum_{j=1}^N e^{-i\mathbf{k} \cdot \mathbf{r}_j}$ is the density operator, and ψ_0 is the many-body ground state (which is, in fact, unknown for ^4He). Note that this trial state automatically builds in the favorable correlations of the ground state. The energy of this excited state, $\Delta(\mathbf{k}) = \langle \phi_{\mathbf{k}} | H - E_0 | \phi_{\mathbf{k}} \rangle / \langle \phi_{\mathbf{k}} | \phi_{\mathbf{k}} \rangle$, can be simply evaluated:

$$\Delta(\mathbf{k}) = \frac{N^{-1} \langle \psi_0 | \rho_{\mathbf{k}}^\dagger [H, \rho_{\mathbf{k}}] | \psi_0 \rangle}{N^{-1} \langle \psi_0 | \rho_{\mathbf{k}}^\dagger \rho_{\mathbf{k}} | \psi_0 \rangle} \equiv \frac{f(\mathbf{k})}{S(\mathbf{k})}, \quad (5)$$

In the last term the numerator is the “oscillator strength” and takes on the universal value $f(\mathbf{k}) = \hbar^2 k^2 / 2m$, and $S(\mathbf{k})$ is the static structure factor, which is directly measurable by means of neutron scattering (it is here, in fact, where the He-He correlations of the ground state are “encoded”). Using the experimental results for $S(\mathbf{k})$, Feynman could calculate a spectrum of remarkable quality, showing the phonon-like spectrum at small wavevectors, and a roton minimum at wavevectors comparable to the inverse interatomic distance.²¹

III. THE SINGLE MODE APPROXIMATION IN THE QUANTUM HALL EFFECT

The applicability of the SMA to *fermion* systems is also well established for two- and three-dimensional systems in the absence of magnetic fields, giving a good approximation for the plasmons at long wavelength. For 2DES in presence of a magnetic field it correctly gives the zero-wavevector magnetoplasmon at $\omega_c = eB/m_e$, a result that is guaranteed by Kohn’s theorem,²⁴ which states that the dipolar excitation is saturated by the cyclotron mode (this results in the modes of interest—the *intra-LL* excitations—having *quadrupolar* character, i.e. with an oscillator strength $\bar{f} \propto q^4$).

For excitations fully contained within a single LL, the cyclotron mode is not of primary interest. In 1985 Girvin, MacDonald and Platzman (GMP) proposed an ingenious ansatz for *projected* excited states:²²




FIG. 2: (Color online) Pair correlation function $g(\mathbf{r})$ for a $\nu = 1/3$ nematic, a $\nu = 1/5$ tetratic, and a $\nu = 1/7$ hexatic. In all cases $g(\mathbf{r}) = \mathcal{O}[r^{2/\nu}]$ for small r and $g(\infty) \rightarrow 1$. The fact that $g(\mathbf{r}) \simeq 1$ over a large area is a guarantee that the calculations are able to reproduce the thermodynamic limit.

$$|\psi_{\mathbf{q}}\rangle = \bar{\rho}_{\mathbf{q}} |\psi_0\rangle, \quad (6)$$

where $\bar{\rho}_{\mathbf{q}}$ is the *projected* density operator:²⁵

$$\begin{aligned} \bar{\rho}_{\mathbf{q}} &= \sum_{m,m'} \langle 0, m' | e^{-i\mathbf{q}\cdot\mathbf{r}} | 0, m \rangle a_{0,m}^\dagger a_{0,m} \\ &= \sum_{j=1}^N \overline{e^{-i\mathbf{q}\cdot\mathbf{r}}} = \sum_{j=1}^N e^{-|\mathbf{q}|^2/2} e^{-iq^* z_j/2} e^{-iq \frac{\partial}{\partial z_j}}, \end{aligned} \quad (7)$$

where $|0, m\rangle$ correspond to single-particle states in the lowest LL and angular momentum m , and $a_{0,m}^\dagger$ is the particle creator operator for such state. As in Feynman's ansatz [Eq. (4)], Eq. (6) preserves the favorable correlations of the ground state. The exclusion of inter-LL excitations eliminates the problem with the saturation of the dipolar mode. The excited states have a compelling description, in first quantized form:²⁵

$$\begin{aligned} \bar{\rho}_{\mathbf{q}} \psi(z_1, \dots, z_N) &= \sum_{j=1}^N e^{-|\mathbf{q}|^2/2} e^{-iq^* z_j/2} \times \\ &\times \psi(z_1, \dots, z_{j-1}, z_j - iq, z_{j+1}, \dots, z_N), \end{aligned} \quad (8)$$

which corresponds to shifting each electron by $\hat{e}_z \times \mathbf{q}$ and superimposing these N configurations with an amplitude $e^{-iq^* z_j/2}$.

Similarly to Eq. (5), the excitation spectrum can be readily obtained

$$\bar{\Delta}_{\mathbf{q}} = \frac{(2N)^{-1} \langle \psi_0 | [\bar{\rho}_{\mathbf{q}}^\dagger, [\bar{H}, \bar{\rho}_{\mathbf{q}}]] | \psi_0 \rangle}{N^{-1} \langle \psi_0 | \bar{\rho}_{\mathbf{q}}^\dagger \bar{\rho}_{\mathbf{q}} | \psi_0 \rangle} \equiv \frac{\bar{f}(\mathbf{q})}{\bar{S}(\mathbf{q})}. \quad (9)$$

The projected oscillator strength comes from the non-commutation of the projected density operator with terms in the *potential* energy part of the Hamiltonian also projected onto the LLL:

$$\bar{H} = \frac{1}{2} \sum_{\mathbf{q}} v_{\mathbf{q}} (\bar{\rho}_{\mathbf{q}}^\dagger \bar{\rho}_{\mathbf{q}} - N e^{-q^2 l^2/2}). \quad (10)$$

Since $[\bar{\rho}_{\mathbf{k}}, \bar{\rho}_{\mathbf{q}}] = (e^{k^* q/2} - e^{k q^*/2}) \bar{\rho}_{\mathbf{k}+\mathbf{q}}$, we find²²

$$\begin{aligned} \bar{f}(\mathbf{q}) &= 2 e^{-|\mathbf{q}|^2/2} \sum_{\mathbf{k}} \sin^2\left(\frac{\mathbf{q} \times \mathbf{k}}{2}\right) \bar{S}(\mathbf{k}) \times \\ &\times [v_{\mathbf{k}-\mathbf{q}} e^{\mathbf{k}\cdot\mathbf{q}} e^{-|\mathbf{q}|^2/2} - v_{\mathbf{k}}], \end{aligned} \quad (11)$$

For its part, the projected static structure factor $\bar{S}(\mathbf{q})$ can be calculated from:^{22,25}

$$\bar{S}(\mathbf{q}) = S(\mathbf{q}) - (1 - e^{-q^2/2}), \quad (12)$$

where $S(\mathbf{q})$ is the *unprojected* static structure factor:

$$S(\mathbf{q}) - 1 = \rho_0 \int d^2 r e^{-i\mathbf{q}\cdot\mathbf{r}} [g(\mathbf{r}) - 1], \quad (13)$$

the Fourier transform of the pair correlation function

$$g(\mathbf{r}) = n_e^{-2} \langle \sum_{i \neq j}^N \delta(\mathbf{r}_i - \mathbf{r}) \delta(\mathbf{r}_j) \rangle \quad (14)$$

which is obtainable from the ground state via, e.g., Monte Carlo (MC) simulations.^{12,13,22} In our case, we considered BRS states [Eq. (2)] corresponding to a $\nu = 1/3$ nematic, a $\nu = 1/5$ tetratic, and a $\nu = 1/7$ hexatic. Figure 2 depicts the pair correlation function for various states. In all cases, correlation functions and SMA excitation spectra were computed for numerous α 's. The angle dependence significantly increases the burden in the MC simulations since the full angle-dependent $g(\mathbf{r})$ is needed, rather than the considerably simpler angle-averaged $g(r)$ of the isotropic systems. The accurate calculation of $\bar{f}(\mathbf{q})$, with its angular-dependent exponentially large factors required high quality $\bar{S}(\mathbf{q})$ and hence $g(\mathbf{r})$. To put things in perspective, to achieve ca. 1% accuracy in $g(\mathbf{r})$, $\mathcal{O}[10,000]$ counts were accumulated for each 0.01×0.01 (in units of l_0) $\mathbf{r}$ -box in the original histograms (the small boxes were necessary to have precision in the Fourier transform at high wavevectors). This required, for each filling factor ν and anisotropy generating α , to run approximately $4\text{--}8 \times 10^7$ MC steps in systems of $N_e = 200\text{--}400$ electrons taking 100–200 cpu $\times$ days of computation for each (ν, α) in a 2 GHz Athlon cluster (see Ref. [13] for further details). A relatively large N_e is required so that the simulations are able to reproduce a system in the thermodynamic limit. The fact that $g(\mathbf{r}) \simeq 1$ over a large area is a guarantee of that achievement (Fig. 2).

Figure 3 shows the *projected* static structure factors $\bar{S}(\mathbf{q})$ for a $\nu = 1/3$ nematic, a $\nu = 1/5$ tetratic, and a $\nu = 1/7$ hexatic. From $\bar{S}(\mathbf{q})$, the oscillator strength $\bar{f}(\mathbf{q})$ is computed using Eq. (11). Analysis of $\bar{S}(\mathbf{q})$ and $\bar{f}(\mathbf{q})$ shows that *both* are $\mathcal{O}[q^4]$ for small $\mathbf{q}$.²² This restriction on the small wavevector behavior originates in Kohn's theorem,²⁴ as all of the $\mathcal{O}[q^2]$




FIG. 3: (Color online) Projected static structure factors $\bar{S}(\mathbf{q})$ for a $\nu = 1/3$ nematic, a $\nu = 1/5$ tetratic, and a $\nu = 1/7$ hexatic. In all cases, $\bar{S}(\mathbf{q}) = \mathcal{O}[q^4]$ for small q , and $\bar{S}(\infty) \rightarrow 0$.

pieces in the *unprojected* parts are saturated by the (uninteresting) inter-LL excitations at $\hbar\omega_c$.

Figure 4 presents some of our results for the excitation spectra in the lowest LL. The results are consistent with those obtained by GMP for the isotropic $\nu = 1/3$ and $1/5$ FQHE cases²² which were qualitatively confirmed experimentally.²⁶ They show that the collective excitation spectrum remains gapped, albeit with a deep magnetoroton, for modes coupled to the ground state via the density operator. Not surprisingly, BRS states have significant anisotropy in their spectra. However, it is noteworthy that for the nematic and tetratic cases the spectrum is *singular*, with an angle dependence on the excitation energy $\Delta(\mathbf{q})$ as $\mathbf{q} \rightarrow 0$. By contrast, the hexatic liquid crystal has a regular spectrum in the long wavelength limit. The apparent disparity have, of course, to do with the different rotational symmetries of the different states: as $\bar{\Delta} = \bar{f}/\bar{S}$, and both numerator and denominator are $\mathcal{O}[q^4]$ for small $\mathbf{q}$, there is no possibility of generating a C_6 symmetric form with terms that can only depend on q_x^4 , q_y^4 , and $q_x^2 q_y^2$. This can also be understood from the point of view of an effective elasticity theory for 2DES (valid in the long wavelength limit, $q \rightarrow 0$):²⁷ the elasticity tensor being of the fourth rank is not compatible with a six-fold rotational symmetry. This elastic interpretation will be published elsewhere.¹⁷

The presence of this singular spectrum suggests that microwave conductivity experiments^{8,26} may be able to discern such structures as a signature of, e.g., the quantum Hall nematic suggested by magnetotransport experiments.^{5,6}

Another interesting feature (Fig. 5) is that the magnetoroton minima and the gap at the origin appears to collapse at some finite anisotropy factor α . The appearance of elastic modes that go soft may be a precursor of the appearance of charge density waves in the system.¹³ Spectra in higher LL's can be obtained by multiplying the projected density operator $\bar{\rho}_{\mathbf{q}}$ by $L_L(q^2/2)$, where $L_L(x)$ is the Laguerre polynomial and L corresponds to the desired LL.²⁸ The rightmost panel of Fig. 4 shows the modification of the SMA spectrum for the first excited LL. Numerical error due to the large $L_L(q^2/2)$ factors

makes it difficult to get reliable results for higher LL's at this point.

Theoretical predictions of which type of order is lowest in *free energy* (required to decide which state is favorable at *finite temperatures*, as shown in Refs. [6,7]) are still incomplete, however, as the entropy is likely to be dominated by gapless modes originating, e.g., from the Goldstone modes associated with the spontaneous breaking of the continuous rotational symmetry of the isotropic states. Generalization of the SMA to gapless rotational modes (see e.g. 15) will require, however, 3-body operators which demanding considerably higher computing capabilities.²⁹

IV. SUMMARY

Summarizing: in connection with the recent variety of experimental evidence supporting liquid crystalline phases in quantum Hall systems,^{5,6,7,8} we have calculated the collective excitation spectrum in the SMA approximation for a quantum Hall liquid crystal. We found that the spectrum of excitations coupled to the ground state by the density operator remains gapped, but develops a significant anisotropy, which in the case of the nematic and tetratic liquid crystals has a singular gap in the long wavelength limit.

V. ACKNOWLEDGMENTS

We would like to acknowledge helpful discussions with G. Vignale, O. Ciftja, A.T. Dorsey, M. Fogler, A. MacDonald, S. Girvin, E. Fradkin, H. Fertig, and J. Jain. Acknowledgment is made to the University of Missouri Research Board and Research Council, and to the Donors of the Petroleum Research Fund, administered by the American Chemical Society, for support of this research.

¹ K. v.Klitzing, G. Dorda and M. Pepper, Phys. Rev. Lett. **45**, 494 (1980); D.C. Tsui, H.L. Stormer and A.C. Gossard, Phys. Rev. Lett. **48**, 1559 (1982).

² R.B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).

³ For reviews see: *Perspectives in quantum Hall effects*, ed. by S. Das Sarma and A. Pinczuk (Wiley, New York 1996); *The quan-*




FIG. 4: (Color online) Single mode approximation spectra for a $\nu = 1/3$ nematic, a $\nu = 1/5$ tetratic, and a $\nu = 1/7$ hexatic. Note the dramatic angular dependence of the spectra, and the appearance of a singular gap as $q \rightarrow 0$ for the nematic and tetratic. The lower right panel shows also the spectrum for the $\nu = 1/3$ nematic in the first excited LL ($L = 1$).




FIG. 5: Possible collapse of the magnetoroton minimum as function of the anisotropy parameter α . Dotted lines are guides to the eye.

tum Hall effect, 2nd Ed., ed. by R.E. Prange and S.M. Girvin (Springer-Verlag, New York 1990).

- ⁴ J. Jain, Phys. Today **53**, 39 (2000); *Composite Fermions*, ed. by O. Heinonen (World Scientific, New York, 1998).
- ⁵ M.P. Lilly, K.B. Cooper, J.P. Eisenstein, L.N. Pfeiffer and K.W. West, Phys. Rev. Lett. **82**, 394, (1999); R.R. Du, D.C. Tsui, H.L. Stormer, L.N. Pfeiffer, K.W. Baldwin and K.W. West, Solid State Comm. **109**, 389 (1999).
- ⁶ G. Gervais, L.W. Engel, H.L. Stormer, D.C. Tsui, K.W. Baldwin, K.W. West, and L.N. Pfeiffer, Phys. Rev. Lett. **93**, 266804 (2004).
- ⁷ W. Pan, H.L. Stormer, D.C. Tsui, L.N. Pfeiffer, K.W. Baldwin, and K.W. West, Phys. Rev. Lett. **88**, 176802 (2002).
- ⁸ R.M. Lewis, Y. Chen, L.W. Engel, D.C. Tsui, P.D. Ye, L.N. Pfeiffer, and K.W. West Phys. Rev. Lett. **93**, 176808 (2004).
- ⁹ M. M. Fogler, A.A. Koulakov and B.I. Shklovskii, Phys. Rev. B **54**, 1853 (1996); R. Moessner and J. T. Chalker, Phys. Rev. B **54**, 5006 (1996).
- ¹⁰ E. Fradkin and S.A. Kivelson, Phys. Rev. B **59**, 8065 (1999); S.A. Kivelson, E. Fradkin, and V.J. Emery, Nature (London) **393**, 550 (1998).
- ¹¹ C. Wexler and A.T. Dorsey, Phys. Rev. B **64**, 115312 (2001).

- ¹² O. Ciftja and C. Wexler, Phys. Rev. B **65**, 045306 (2002); *ibid*, 205307 (2002); A.J. Schmidt, O. Ciftja, and C. Wexler, *ibid* **67**, 155315 (2003). C. Wexler and O. Ciftja, J. Phys.: Condens. Matter **14**, 3705 (2002).
- ¹³ O. Ciftja, C.M. Lapilli and C. Wexler, Phys. Rev. B **69**, 125320 (2004).
- ¹⁴ A.H. MacDonald and M.P.A. Fisher, Phys. Rev. B **61**, 5724 (2000); F. von Oppen, B. I. Halperin, and A. Stern, Phys. Rev. Lett. **84**, 2937 (1999); S.-Y. Lee, V.W. Scarola, and J.K. Jain, Phys. Rev. Lett. **87**, 256803 (2001).
- ¹⁵ L. Radzihovsky and A.T. Dorsey, Phys. Rev. Lett. **88**, 216802 (2002); M.M. Fogler, Europhys. Lett. **66**, 572 (2004).
- ¹⁶ C.M. Lapilli and C. Wexler, in *ICPS27* ed. by J. Menendez and C. van de Walle (AIP Conf. Proc., New York, 2005); C. Wexler and C.M. Lapilli, in *Condensed Matter Theories*, vol. 20, ed. by R.M. Panoff and J.W. Clark (Nova Science Publishers, 2005).
- ¹⁷ C.M. Lapilli, C. Wexler, and G. Vignale, Bull. Am. Phys. Soc. II Vol. 50 (1), 1272 (2005), paper in preparation.
- ¹⁸ K. Musaelian and R. Joynt, J. Phys.: Condens. Matter **8**, L105 (1996); L. Balents, Europhys. Lett. **33**, 291 (1996).
- ¹⁹ J.M. Kosterlitz and D.J. Thouless, J. Phys. C **6**, 1181 (1973); J.M. Kosterlitz, *ibid*. **7**, 1046 (1974).
- ²⁰ B.I. Halperin and D.R. Nelson, Phys. Rev. Lett. **41**, 121 (1978); A.P. Young, Phys. Rev. B **19**, 1855 (1979).
- ²¹ R.P. Feynman, Phys. Rev. **91**, 1291 (1953); *ibid*. **94**, 262 (1954).
- ²² S.M. Girvin, A.H. MacDonald, and P.M. Platzman, Phys. Rev. Lett. **54**, 581 (1985); Phys. Rev. B **33**, 2481 (1986).
- ²³ K. Park and J.K. Jain, Solid State Comm. **115**, 353 (2000).
- ²⁴ W. Kohn, Phys. Rev. **123**, 1242 (1961).
- ²⁵ S.M. Girvin, Phys. Rev. B **30**, 558 (1984).
- ²⁶ A. Pinczuk, B.S. Dennis, L.N. Pfeiffer, and K. West, Phys. Rev. Lett. **70**, 3983 (1993); Physica B **249**, 40 (1998).
- ²⁷ S. Conti and G. Vignale, J. Cond. Mat. Phys. **10**, L779 (1998).
- ²⁸ See, e.g., A.H. MacDonald and S.M. Girvin, Phys. Rev. B **33**, 4009 (1986); and Refs. [12,13].
- ²⁹ M.M. Fogler, private communication.